



UPDATE:

Update to communications issued 25th April 2024
Material updates are shown in bold

Medicine Supply Notification

MSN/2024/047U

Daktacort[®] (hydrocortisone 1% / miconazole 2%) cream

Tier 2 – medium impact*

Date of issue: 08/05/2024

Link: [Medicines Supply Tool](#)

Summary

- Daktacort[®] (hydrocortisone 1% / miconazole 2%) cream (30g, POM) has been discontinued with remaining supplies due to be exhausted by the end of May 2024.*
- Daktacort Hydrocortisone[®] (hydrocortisone 1% / miconazole 2%) cream 15g (OTC) remains available but cannot support increased demand until October 2024.**
- Alternative cream formulations containing a different antifungal in combination with hydrocortisone (0.5% or 1%) remain available and can support increased demand.

Actions Required

Clinicians should:

- not initiate any new patients on Daktacort[®] cream (30g);
- consider prescribing an alternative antifungal- hydrocortisone combination cream, ensuring that the patient is not intolerant to any of the excipients (see Supporting information); and
- consider prescribing Daktacort Hydrocortisone cream (15g) if it is deemed the most appropriate product when it is able to support increased demand in October 2024, noting use may be off label in light of its more limited range of licensed indications (see Supporting information).

*Classification of Tiers can be found at the following link:

<https://www.england.nhs.uk/publication/a-guide-to-managing-medicines-supply-and-shortages/>

Supporting information

Table 1: Antifungal- hydrocortisone combination creams

Product	Constituents	indication
Daktacort [®] (POM)*	Miconazole nitrate 2%w/w Hydrocortisone 1%w/w. <u>Excipients with known effect:</u> 2 mg/g benzoic acid (E210) & 0.052 mg/g butylated hydroxyanisole (E320)	<ul style="list-style-type: none"> Inflamed dermatoses where infection by susceptible organisms and inflammation co-exist, e.g. intertrigo and infected eczema. Moist or dry eczema or dermatitis including atopic eczema, primary irritant or contact allergic eczema or seborrhoeic eczema including that associated with acne. Intertriginous eczema including inflammatory intertrigo, perianal and genital dermatitis. <p>Organisms susceptible to miconazole are dermatophytes and pathogenic yeasts (e.g. Candida spp.). Also, many Gram-positive bacteria including most strains of Streptococcus and Staphylococcus.</p>
Daktacort Hydrocortisone [®] (P)**	Miconazole nitrate 2% w/w; hydrocortisone 1% w/w <u>Excipients with known effect:</u> Benzoic acid (E210) 2 mg/g & butylhydroxyanisole (E320) 0.052 mg/g	Athlete's foot and candidal intertrigo where there are co-existing symptoms of inflammation.
Canesten HC [®]	Each gram of cream contains:10mg clotrimazole & 10 mg hydrocortisone (1%) <u>Excipients with known effect:</u> Cetostearyl alcohol 100mg & Benzyl alcohol 20mg (in each gram of cream)	<ul style="list-style-type: none"> Skin infections where co-existing symptoms of inflammation, e.g. itching, require rapid relief: Dermatomycoses due to dermatophytes (e.g. Trichophyton species), moulds and other fungi & due to yeasts (Candida species) and skin diseases showing secondary infection with these fungi. Nappy rash where infection due to Candida albicans is present. Candidal vulvitis, candidal balanitis and candidal intertrigo.
Timodine [®]	Nystatin 3.00 %w/w Dimeticone 350 10.00%w/w Hydrocortisone 0.50%w/w Benzalkonium chloride solution 0.20%w/w <u>Excipients with known effect:</u> Cetostearyl alcohol 1.51%w/w Butylated hydroxyanisole compound 0.40%w/w [containing butylated hydroxyanisole (E320) 20%, propyl gallate 10%, citric acid 10% and 60% propylene glycol (E1520)] Methyl hydroxybenzoate (E218) 0.10 w/v Propyl hydroxybenzoate (E216) 0.10% w/w Sorbic acid 0.10% w/w	<ul style="list-style-type: none"> Dermatoses, including intertrigo, eczema, seborrhoeic dermatitis, hand dermatitis, and pruritis ani et vulvae, in which infection with Candida albicans is a factor. Severe nappy rash in which infection with Candida albicans is a factor.
Nystaform HC [®]	Nystatin 100,000 units/g Chlorhexidine 1%w/w Hydrocortisone 0.5%w/w <u>Excipients with known effect:</u> Cetostearyl alcohol 100 mg & Benzyl alcohol 10 mg (per gram of cream)	Infected dermatoses where fungal (particularly monilial) and/or bacterial infections are present.

[Links to further information](#)

[SmPC for Daktacort® cream](#)

[SmPC for Daktacort Hydrocortisone® cream](#)

[SmPC for Canesten HC® cream](#)

[BNF: Hydrocortisone with miconazole](#)

[BNF: Hydrocortisone with clotrimazole](#)

[BNF: Hydrocortisone with chlorhexidine hydrochloride and nystatin](#)

[BNF: Hydrocortisone with benzalkonium chloride, dimeticone and nystatin](#)

[SmPC for Timodine® cream](#)

[SmPC for Nystaform HC®](#)

Enquiries

If you have any queries, please contact DHSCmedicinesupplyteam@dhsc.gov.uk.